

Host-Derived Sialic Acids Are an Important Nutrient Source Required for Optimal Bacterial Fitness *In Vivo*

Nathan D. McDonald, Jean-Bernard Lubin, Nityananda Chowdhury, E. Fidelma Boyd

Department of Biological Sciences, University of Delaware, Newark, Delaware, USA

ABSTRACT A major challenge facing bacterial intestinal pathogens is competition for nutrient sources with the host microbiota. *Vibrio cholerae* is an intestinal pathogen that causes cholera, which affects millions each year; however, our knowledge of its nutritional requirements in the intestinal milieu is limited. In this study, we demonstrated that *V. cholerae* can grow efficiently on intestinal mucus and its component sialic acids and that a tripartite ATP-independent periplasmic SiaPQM strain, transporter-deficient mutant NC1777, was attenuated for colonization using a streptomycin-pretreated adult mouse model. In *in vivo* competition assays, NC1777 was significantly outcompeted for up to 3 days postinfection. NC1777 was also significantly outcompeted in *in vitro* competition assays in M9 minimal medium supplemented with intestinal mucus, indicating that sialic acid uptake is essential for fitness. Phylogenetic analyses demonstrated that the ability to utilize sialic acid was distributed among 452 bacterial species from eight phyla. The majority of species belonged to four phyla, *Actinobacteria* (members of *Actinobacillus*, *Corynebacterium*, *Mycoplasma*, and *Streptomyces*), *Bacteroidetes* (mainly *Bacteroides*, *Capnocytophaga*, and *Prevotella*), *Firmicutes* (members of *Streptococcus*, *Staphylococcus*, *Clostridium*, and *Lactobacillus*), and *Proteobacteria* (including *Escherichia*, *Shigella*, *Salmonella*, *Citrobacter*, *Haemophilus*, *Klebsiella*, *Pasteurella*, *Photobacterium*, *Vibrio*, and *Yersinia* species), mostly commensals and/or pathogens. Overall, our data demonstrate that the ability to take up host-derived sugars and sialic acid specifically allows *V. cholerae* a competitive advantage in intestinal colonization and that this is a trait that is sporadic in its occurrence and phylogenetic distribution and ancestral in some genera but horizontally acquired in others.

IMPORTANCE Sialic acids are nine carbon amino sugars that are abundant on all mucous surfaces. The deadly human pathogen *Vibrio cholerae* contains the genes required for scavenging, transport, and catabolism of sialic acid. We determined that the *V. cholerae* SiaPQM transporter is essential for sialic acid transport and that this trait allows the bacterium to outcompete non-catabolizers *in vivo*. We also showed that the ability to take up and catabolize sialic acid is prevalent among both commensals and pathogens that colonize the oral cavity and the respiratory, intestinal, and urogenital tracts. Phylogenetic analysis determined that the sialic acid catabolism phenotype is ancestral in some genera such as *Yersinia*, *Streptococcus*, and *Staphylococcus* and is acquired by horizontal gene transfer in others such as *Vibrio*, *Aeromonas*, and *Klebsiella*. The data demonstrate that this trait has evolved multiple times in different lineages, indicating the importance of specialized metabolism to niche expansion.

Received 24 December 2015 Accepted 25 February 2016 Published 12 April 2016

Citation McDonald ND, Lubin J-B, Chowdhury N, Boyd EF. 2016. Host-derived sialic acids are an important nutrient source required for optimal bacterial fitness *in vivo*. mBio 7(2):e02237-15. doi:10.1128/mBio.02237-15.

Editor Vanessa Sperandio, UT Southwestern Med Center Dallas

Copyright © 2016 McDonald et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported license](#), which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

Address correspondence to E. Fidelma Boyd, fboyd@udel.edu.

Vibrio cholerae is the causative agent of cholera, the deadly diarrheal disease that affects millions each year. To date, there have been seven cholera pandemics, the latest of which, the 7th pandemic, began in 1961 (1–3). Pandemic cholera-causing isolates of *V. cholerae* belong to the O1 serogroup, which is divided into two biotypes: the classical biotype and the El Tor biotype. The current 7th pandemic is of the El Tor biotype, and the classical biotype strains are believed to have caused the first six pandemics of cholera but have now disappeared from the regions of cholera endemicity (1, 3). Additionally, in the early 1990s, a new serogroup, O139, emerged as a leading cause of cholera in what was believed at the time to be the beginning of the 8th pandemic; however, within a few years, the El Tor biotype reemerged as the predominant cause of cholera (1, 3, 4). Interestingly, O139 isolates lack the sialic acid catabolism (SAC) gene cluster (5). Others have

speculated that one explanation of why the El Tor biotype was able to dominate the classical biotype is that it was a consequence of enhanced metabolic fitness (6, 7). They proposed that the ability to produce a neutral fermentation end product rather than an acid by-product from metabolized sugars improves fitness in a range of environments.

With nearly 100 trillion commensal bacteria colonizing the intestine, being highly efficient at scavenging for nutrients becomes essential for survival (8–10). Enhanced metabolic fitness is especially important for gastroenteritis-causing intestinal pathogens, as these pathogens compete for nutrients with the already established microbiota (11–15). The gastrointestinal (GI) tract is lined with a mucus layer serving as a barrier between epithelial cells and bacteria as well as protecting the epithelium from digestive enzymes found in the lumen (16). The main components of

mucus are mucins, which comprise a family of heavily glycosylated proteins that are secreted from the epithelial tissue and form the gel-like mucus layer that covers the cells (16). The glycosylation of mucin is of particular importance for the gut microbiota, as many commensal as well as pathogenic species are able to cleave sugars from mucin in order to use them as nutrients (17, 18). Some of the main sugars that are typically found in mucus include ribose, mannose, hexuronates, galactose, fucose, arabinose, *N*-acetylglucosamine (GlcNAc), and the sialic acid *N*-acetylneuraminic acid (Neu5Ac) (19, 20). Sialic acid is of particular interest because it is the most common terminal carbohydrate present on N- and O-linked glycans, which are predominant in the mucus-covered oral, lung, intestinal, and vaginal surfaces. Additionally, there have been over 50 structural sialic acid derivatives that been characterized, the most abundant of which is Neu5Ac.

Cholera-causing strains of *V. cholerae* carry genes necessary for transport (*siaPQM*) and catabolism (*nanA*, *nanEK*) of sialic acid as well as a gene encoding a scavenging enzyme, sialidase (*nanH*), which can cleave the sialic acid residues from the glycan, allowing them to be taken up into the bacterial cell (5). These genes are contained within the 57-kb *Vibrio* pathogenicity island 2 (VPI-2) region that integrates at a tRNA-serine locus (5). The canonical VPI-2 region is present only in *V. cholerae* O1 serogroup strains; however, the majority of the region has been deleted from O139 serogroup isolates (5, 21). Previously, we have shown that *V. cholerae* N16961, an O1 serogroup El Tor isolate, can utilize Neu5Ac as a sole carbon source and that a mutant lacking the sialic acid aldolase enzyme *N*-acetylneuraminic acid lyase (*NanA*) is defective in colonization of an infant mouse model compared to wild-type *V. cholerae* (22). Additionally, it was demonstrated that *SiaPQM*, the tripartite ATP-independent periplasmic (TRAP) transporter within VPI-2, is solely responsible for the transport of the Neu5Ac sialic acid (23). Four diverse families of sialic acid transporters have been shown to be genetically linked with SAC genes among bacteria: the TRAP transporter system, the type present in most *Vibrio* species; the major facilitator superfamily (MFS)-type NanT transporters; an ATP-binding cassette (ABC)-type transporter; and the sodium solute symporter (SSS)-type transporters (24–28).

In this study, we investigated the ability of *V. cholerae* to grow in mouse intestinal mucus as well as in individual sugar components of mucus as sole carbon sources. We investigated the ability of *V. cholerae* to utilize sialic acid derivatives *N*-glycolylneuraminic acid (Neu5Gc) and 2-keto-3-deoxy-D-glycero-D-galacto-nononic acid (KDN) as sole carbon sources. We determined the significance of sialic acid transport *in vivo* using a streptomycin (Sm)-pretreated adult mouse model of colonization as well as competition persistence assays between the wild-type strain and NC1777, a *SiaPQM* transporter-deficient mutant strain. The results of the *in vivo* study as well as of *in vitro* competition assays suggest that sialic acids are important nutrients that *V. cholerae* scavenges for during host colonization. We determined the distribution of the ability to take up and catabolize sialic acid and mapped the different types of sialic acid transporters onto the phylogeny of the *NanA* protein among bacteria to demonstrate that this phenotype evolved multiple times.

RESULTS AND DISCUSSION

***Vibrio cholerae* utilizes intestinal mucus and its components as sole carbon sources.** During colonization in the GI tract, one of the main sources of nutrients is mucus secreted from the epithelial cells lining the lumen (16). Here, we assessed the ability of *V. cholerae* N16961 to grow in M9 minimal medium supplemented with intestinal mucus (M9M) as the sole carbon source. We demonstrated that *V. cholerae* is able to grow in intestinal mucus, showing no observable lag phase (Fig. 1A). We next determined the growth patterns on the main monosaccharides that are commonly found as part of intestinal mucus: ribose, mannose, hexuronates (gluconate and glucuronate), galactose, fucose, arabinose, *N*-acetylglucosamine (GlcNAc), and Neu5Ac. We found that *V. cholerae* N16961 achieved comparable levels of biomass (maximum optical density [OD]) when grown on glucose, ribose, mannose, gluconate, GlcNAc, or Neu5Ac as the sole carbon source. Only moderate growth was shown with galactose as a sole carbon source (Fig. 1B).

It is unknown whether *V. cholerae* can utilize additional sialic acids such as Neu5Gc, which is a major component of all mammalian mucins, with the exception of mucins of humans, or KDN, a recently identified sialic acid of lower-order vertebrates (29, 30). Our growth analysis showed that N16961 can utilize Neu5Ac and Neu5Gc but not KDN as a sole carbon source (Fig. 2A to C). To determine whether the *SiaPQM* TRAP transporter was required for Neu5Gc transport, we examined a *SiaPQM*-deficient strain, NC1777, for Neu5Gc uptake and found that the mutant did not grow on either Neu5Ac or Neu5Gc (Fig. 2A and B). Interestingly, the growth pattern of *V. cholerae* showed a shorter lag phase on Neu5Ac and Neu5Gc than that seen with *Escherichia coli*, which had an extended lag phase. This suggests that *V. cholerae* *SiaPQM* transports both of these substrates with greater efficiency than *E. coli* BW25113, which is consistent with TRAP transporters being high-affinity transporters for their respective solutes. To further examine the ability of *Vibrio* species to utilize KDN and to ensure that our phenotype was not strain specific, we examined the growth patterns of *Vibrio vulnificus*, a species distantly related to *V. cholerae* and containing an identical sialic acid transport and catabolism region (26, 31). Similarly to what we showed for *V. cholerae*, *V. vulnificus* CMCP6 can utilize Neu5Ac and Neu5Gc but not KDN as a sole carbon source (see Fig. S1 in the supplemental material). To determine whether *SiaPQM* was required for Neu5Gc transport, we examined a *SiaPQM*-deficient strain, JJK0731, for Neu5Gc uptake and also demonstrated that the mutant cannot grow on either Neu5Ac or Neu5Gc (see Fig. S1).

A recent study by Hopkins and colleagues demonstrated the ability of *E. coli* to catabolize both Neu5Gc and KDN (32). In their study, they found that the *E. coli* permease NanT was required for transport of these substrates. They also demonstrated that complementation of a NanT mutant with the *SiaPQM* transporter from *Haemophilus influenzae* also resulted in uptake of both Neu5Gc and KDN but that the affinity for KDN was 50-fold lower than for Neu5Ac (32). We complemented an *E. coli nanT* mutant with *siaPQM*, strain NDM3193C, and showed that in the *E. coli* background, both Neu5Ac and Neu5Gc were transported but not KDN, suggesting that *SiaPQM* is not a transporter for KDN (Fig. 2D). In addition, we also complemented the *V. cholerae* NC1777 mutant with *siaPQM* and showed that Neu5Ac and




FIG 1 Growth analysis of *V. cholerae* N16961 on mouse intestinal mucus and mucus sugars. (A) *V. cholerae* growth in M9 minimal medium supplemented with mucus (M9M). (B) Maximum density (OD₅₉₅) of *V. cholerae* grown in M9 supplemented with glucose, *N*-acetylglucosamine (GlcNAc), galactose, gluconate, mannose, ribose, or Neu5Ac. (C) *V. cholerae* growth in LB media. Data are presented as averages of results of two biological replicates with three technical replicates. Error bars represent standard errors of the means (SEM).

Neu5Gc were transported and catabolized in the complemented mutant (see Fig. S1 in the supplemental material).

The ability to transport sialic acid is essential for *V. cholerae* fitness *in vivo*. To address whether the ability to take up sialic acids is a significant attribute of *V. cholerae in vivo*, we performed *in vivo* competition and persistence assays using a *V. cholerae* streptomycin-pretreated adult mouse model of colonization (33).

Conventionally reared mice contain diverse microbiotas of more than 500 species predominantly from two phyla, *Firmicutes* and *Bacteroidetes*, that prevent infection by many enteric pathogens (reviewed in reference 12). Pretreatment of mice with streptomycin to allow pathogen colonization has been utilized to study the mechanisms of pathogenesis as well as the host response to these pathogens in many studies (34–42). The antibiotic treatment does not change the overall number of bacteria but favors expansion of *Proteobacteria* at the expense of *Bacteroidetes*, with the total number of *Firmicutes* remaining the same (12, 43). Several studies have used this model to examine the role of metabolism, and some have suggested that the initial colonization of a pathogen is dependent on nonlimiting nutrients that are available due to the removal of facultative anaerobes; however, during persistence in the gut, the pathogen must compete for the limiting nutrients (13–15, 44–50). For these reasons, the streptomycin-treated mouse model was used to investigate the competition between wild-type *V. cholerae* and a knock-out strain that cannot transport sialic acid. To accomplish this, we compared NC1777 ($\Delta siaM$) and a β -galactosidase N16961 mutant strain, SAM2338 ($\Delta lacZ$), which allows a blue/white colony screen. First, to confirm that NC1777 and SAM2338 grow similarly to wild-type N16961, we determined growth in LB and M9M at 37°C overnight and found no difference in growth patterns between the two strains (Fig. 1). In addition, we performed *in vitro* competition assays between SAM2338 and N16961 grown overnight in LB at 37°C and found no difference in the CFU counts between the strains (data not shown). Streptomycin-pretreated mice were inoculated with a 1:1 ratio mix of equal amounts of SAM2338 and NC1777, resulting in a final concentration of 1×10^{10} CFU/ml. Following inoculation, at 6, 12, 24, 48, and 72 h postinfection, fecal pellets were collected from each of the mice, the CFU were enumerated, and the competitive index (CI) was calculated. First, we compared SAM2338 and N16961 in coculture *in vivo* and confirmed that, similarly to the *in vitro* assays, there was no competition between the strains, with a CI not significantly different from 1 (data not shown). Then, we examined a coculture of SAM2338 and NC1777 ($\Delta siaM$) and determined that the $\Delta siaM$ strain had a CI close to 1 after 6 h postinfection. However, at 12 h postinfection, the $\Delta siaM$ strain was outcompeted 1.7-fold ($P < 0.001$) by SAM2338 (mean CI of 0.58) (Fig. 3A). This trend continued at time points 24 h, 48 h, and 72 h, with the $\Delta siaM$ strain significantly ($P < 0.001$) outcompeted, with mean CIs of 0.39, 0.37, and 0.33, respectively (Fig. 3A). These data indicate that at the early stages of colonization, sialic acid may not be a limiting nutrient; however, as colonization persists, sialic acid represents a carbon source that strains must compete for and that contributes to fitness. On day 3 of the colonization persistence assay, mice were sacrificed, and the gastrointestinal tracts were removed, sectioned into the small intestine, cecum, and large intestine, homogenized in phosphate-buffered saline (PBS), serially diluted, and plated on LB X-Gal (5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside). It was determined that the $\Delta siaM$ strain was significantly outcompeted in the small intestine, cecum, and large intestine by SAM2338, corroborating the fecal pellet data (Fig. 3B). SAM2338 showed the greatest fitness advantage over the $\Delta siaM$ strain in the small intestine, with a CI of 0.30, followed by the cecum (CI = 0.49) and, finally, the large intestine (CI = 0.59) (Fig. 3B).

Loss of SiaPQM is detrimental *in vitro*. An *in vitro* competition assay was performed using the same inoculum mix for the *in*




FIG 2 Growth analysis of *V. cholerae* N16961 with sialic acid derivatives as sole carbon sources. *V. cholerae* N16961, *V. cholerae* NC1777 ($\Delta siaM$), *E. coli* BW25113, and *E. coli* JW3193 ($\Delta nanT$) in M9 Neu5Ac (A), M9 Neu5Gc (B), or M9 KDN (C) or *E. coli* JW3193 in M9 Neu5Ac, M9 Neu5Gc, or M9 KDN (D) were complemented with *siaPQM* and tested for restored growth on sialic acid derivatives. Growth curves represent results of at least two biological replicates performed in triplicate. Error bars represent SEM.

vivo assays containing equal amounts of NC1777 and SAM2338. The strains were competed against each other in either LB or M9 plus glucose (M9G) (Fig. 4A). The coinoculated cultures were incubated for 24 h, serially diluted, and plated on LB X-Gal. The CI in M9G was 1.06, indicating that neither strain exhibits any growth advantage over the other in this medium; however, in LB, the CI was 0.04 (Fig. 4A). Thus, the *siaM* mutant has a significant defect in competition assays in LB. In order to further investigate the significant defect of the *siaM* mutant cocultured in LB, a time course competition assay was performed in LB and LB supplemented with 20 mM glucose. The CI was calculated at 3, 6, 9, 12, and 24 h postcoinoculation. In LB, at up to 24 h, there was no significant competition between SAM2338 and NC1777; however, at the 24-h time point, the *siaM* mutant was again significantly outcompeted by SAM2338 with a CI that was below the limit of detection of 8.1×10^{-8} CFU/ml (Fig. 4B). When glucose was supplied to the LB media, there was no competition between the strains over the course of the assay (Fig. 4B).

To investigate additional substrates that could require *SiaPQM* for transport, we examined growth of N16961 and NC1777 on 190 different carbon sources, 62 of which were utilized by both strains

under the conditions examined (see Fig. S2 in the supplemental material). There were six substrates in the presence of which NC1777 either showed no growth (Neu5Ac) or reached a lower OD than N16961 (D-glucosamine, L-proline, L-lactic acid, methyl pyruvate, and α -keto-valeric acid). With the exception of D-glucosamine, all of these substrates are structurally similar, being carboxylic acids or carboxylic acid derivatives. Additionally, previous studies have shown evidence of TRAP substrate binding proteins able to bind each of these compounds except D-glucosamine (51, 52). The main components of LB are tryptone and yeast extract; the former is composed of peptides of casein, which contain a high number of proline residues and sialic acid. Our data suggest that *SiaPQM* may play a secondary role in the transport of additional substrates, which may explain the significant defect in *in vitro* competitions assays in LB.

In an attempt to more closely mimic an *in vivo* nutrient environment, strains were competed against one another in mouse mucus isolated from the small intestine, large intestine, and cecum. *In vitro* competition assays in M9M were performed as previously described. The cultures were incubated for 24 h, serially diluted, and plated on LB X-Gal plates. The CI values for the small




FIG 3 *Vibrio cholerae* *in vivo* competition assays. *In vivo* competition assays were performed using streptomycin-pretreated mice which were orally inoculated with an equal amounts of SAM2338 ($\Delta lacZ$) and NC1777 ($\Delta siaM$). (A) Competitive index data were determined by homogenizing fecal pellets collected at 6, 12, 24, 48, and 72 h postinoculation followed by plating on LB X-Gal for a blue/white screen. The competitive index is calculated as follows: $CI = \text{ratio out}_{(NC1777 / SAM2338)} / \text{ratio in}_{(NC1777 / SAM2338)}$. (B) Upon completion of the *in vivo* persistence assay (72 h), mice were sacrificed, gastrointestinal tracts were removed and sectioned into small intestine (SI), large intestine (LI), and cecum (CE), and CFU values were calculated. *In vivo* assays were performed with two biological replicates and $n = 10$ mice. Error bars represent SEM. Statistics were calculated using a one-sample t test with the means compared to the hypothetical CI of 1. Asterisks represent the following: ***, $P < 0.01$.

intestine, large intestine, and cecum were 0.61, 0.58, and 0.67, respectively (Fig. 4A). These results showed that the sialic acid transporter mutant was less fit in each of the three mucus samples and that the ability to transport and catabolize sialic acids is essential for the optimal fitness of cholerae in mucosal environments.

Distribution and phylogenetic analysis of N-acetylneuraminic acid lyase (NanA) among bacteria. Since we and others have shown how significant sialic acid utilization is *in vivo* among a number of pathogens (22, 26, 28), we next determined the prevalence of sialic acid utilization genes in all available completed bacterial genomes. We identified 452 bacterial species with the potential to utilize sialic acid as a carbon source from ~7,400 species representing 23 major bacterial groups, nearly a 10-fold increase




FIG 4 *In vitro* competition assays. Results of assays of *in vitro* competition between *V. cholerae* strain SAM2338 ($\Delta lacZ$) and the sialic acid transporter mutant NC1777 ($\Delta siaM$) are presented. (A) Competition assays were performed in M9G, LB, or M9 supplemented with small-intestine (SI), large-intestine (LI), or cecum mucus. (B) Time course *in vitro* competition assays were performed at 3, 6, 9, 12, and 24 h postinoculation in either LB or LB supplemented with glucose (0.4% [wt/vol]). *In vitro* data represent averages of results of two biological replicates, and error bars represent SEM. Statistics were calculated using an unpaired Student's t test with a 95% confidence interval. All samples were compared to M9G to determine statistical significance. Asterisks represent the following: **, $P < 0.05$.

from our previous study (26). These 452 species were comprised of primarily commensal and pathogen species and encompassed eight phyla: *Actinobacteria*, with 83 representatives; *Bacteroidetes*, with 70 representatives; *Firmicutes*, with 134 representatives; *Fusobacteria*, with 9 representatives; *Planctomycetes*, with 6 representatives; *Proteobacteria*, with 142 representatives; *Spirochaetes*, with 5 representatives; and *Verrucomicrobia*, with 3 representatives. Of the 300 species representing *Crenarchaeota*, *Cyanobacteria*, *Deinococcus-Thermus*, and *Euryarchaeota* from environmental samples, none contained NanA; similarly, of the 188 representatives of the *Bacillus* genus, predominantly soil organisms, only one species had the ability to utilize sialic acid. Within some species, SAC is present in all strains; in other species, it is present in only a subset of strains. The 452 species are comprised of Gram-positive and Gram-negative species, and a few species from divergent phyla are within each of these major groups. These data demonstrate that SAC is phylogenetically widespread but is sporadic in its occurrence within phyla, families, genera, and even




FIG 5 NanA phylogeny. A phylogenetic tree of NanA is shown. The tree was constructed using the neighbor-joining tree method and the p-distance model of evolutionary distances as parameters and MEGA6 (75–77). Letters indicate major bacterial phyla present in the corresponding NanA clade. Colors indicate the major phylogenetic groups that contain *nanA*. Red, *Actinobacteria*; purple, *Bacteroidetes*; lime green, *Firmicutes*; dark green, *Fusobacteria*; dark red, *Planctomycetes*; blue, *Proteobacteria*; yellow, eukaryotes; gray, *Spirochaetes*.

species. The data suggest that the ability to utilize sialic acid as a carbon source was acquired by a select group of species that are colonizers.

In order to examine the evolutionary history of this trait more closely, we constructed a phylogenetic tree of NanA among the 452 species and examined the branching patterns among these species (Fig. 5). In addition, we mapped the different types of transporters associated with the sialic acid catabolism genes from each species onto this tree. For ease of analysis, we have divided the tree into 20 major clusters designated A to T on the tree (Fig. 5). What is immediately apparent from this analysis is that representatives of each of the eight major phyla do not cluster together, indicating different evolutionary histories of NanA among the groups (Fig. 5).

For example, species from the phylum *Proteobacteria* that contain a NanA homologue are found in 6 highly divergent clusters designated A, G, H, I, O, and P, indicating that this trait has evolved several times in this phylum (Fig. 5). Cluster A is a tight group of closely related NanA alleles from the family *Pasteurel-*

laceae represented by *Aggregatibacter*, *Haemophilus*, and *Pasteurella* species, and all encode a TRAP-type sialic acid transporter, with the exception of *H. ducreyi*, which contains an ABC sialic acid transporter (53) (see Fig. S3A in the supplemental material). Sialic acid transport and catabolism was demonstrated in *H. influenzae*, *H. ducreyi*, and *P. multocida* (24, 53–57). Nested within cluster A are species of the phylum *Actinobacteria*, which suggests that NanA in these species may have been acquired from *Proteobacteria* (see Fig. S3A). Branching separately within cluster A are 8 *Fusobacteria* representatives, which all also have a sialic acid TRAP transporter system.

The second *Proteobacteria* group is cluster G and is comprised of *Enterobacteriaceae* species, *E. coli*, *Shigella* spp., *Salmonella* spp., *Citrobacter* spp., and *Klebsiella* spp., all of which form a tight group with short branch lengths and contain species that are known pathogens, and many have been shown to utilize sialic acid as a carbon source (see Fig. S3B in the supplemental material) (25, 26, 58–60).

We identified 31 species of *Aliivibrio*, *Photobacterium*, and

Vibrio that can putatively transport and catabolize sialic acid. Three species, *Vibrio nigripulchritudo*, *V. sinaloensis*, and *V. orientalis*, contained two copies of NanA which share <30% amino acid identity. One NanA copy clustered with all NanA proteins from all other *Vibrio* species (cluster O), whereas the second copy branched in a distant lineage with members of the *Enterobacteriaceae* (cluster I) (see Fig. S3C in the supplemental material). Cluster O is comprised solely of 23 *Vibrio* species, and all contain a TRAP transporter (see Fig. S3D in the supplemental material). Cluster P is comprised mainly of *Aliivibrio*, *Photobacterium*, *Enterobacteria*, *Psychromonas*, *Pseudoalteromonas*, and *Yersinia* species. Subgroup P1 is comprised of all 12 *Yersinia* species, and NanA is present in all strains of each species, indicating that this is an ancestral trait within the genus. Subgroup P2 contains *Plesiomonas shigelloides*, *Aeromonas hydrophila*, and *A. veronii* isolates, human and fish pathogens. Five *A. hydrophila* strains from among the 31 sequenced and 2 *A. veronii* strains from among 12 sequenced contained NanA. The gene orders of the *A. hydrophila* and *A. veronii* SAC clusters differed, and both integrase and transposase genes were associated with both regions, suggesting that the SAC cluster represents a horizontally acquired trait in this genus. Indeed, in *A. hydrophila* strain ML09-119, the sialic acid region is on a genomic island that also contains the genes for fucose utilization, indicating the acquisition of a metabolic island. It is of interest that *A. hydrophila* ML09-119 represents an invasive species from Asia that caused serious disease epidemics in United States catfish farms. The island region is absent from indigenous strains, suggesting that acquisition by the Asian strain may be one of the factors that make them so deadly. These species branched with NanA from three *Photobacterium* species, and all contained an SSS-type transporter. Subgroup P4 represented mainly *Pseudoalteromonas* species, and P5 contained *Klebsiella*, *Edwardsiella*, and *Proteus* species (see Fig. S3D). In *Klebsiella oxytoca*, a second divergent copy of NanA is found within cluster P4. Our growth curve analysis of *Aeromonas hydrophila*, *Enterobacter aerogenes*, *Klebsiella oxytoca*, and *Providencia rettgeri* on Neu5Ac demonstrated that all four species can utilize sialic acid as a sole carbon source (see Fig. S4).

The phylum *Firmicutes* is represented by 5 divergent clusters, and either an SSS transporter (clusters B, C, D, and E) or an ABC transporter (cluster J) is associated with the SAC genes in these groups (Fig. 5). Cluster B contains 13 *Clostridium* species, and within cluster C is a sole representative of the genus *Bacillus*, *B. aquimaris*, along with 3 *Planococcus* species, all of which contain a TRAP transporter, which is highly unusual for Gram-positive bacteria (see Fig. S5A in the supplemental material). Cluster D represents all *Staphylococcus* species, and most *Lactobacillus* species are present in cluster E (see Fig. S5B). NanA appears to be present in all sequenced representatives of *Staphylococcus*, indicating that SAC represents an ancestral trait. Both *Staphylococcus aureus* and *Lactobacillus plantarum* have been shown to utilize sialic acid (61, 62). Cluster J, which is related to clusters B, D, and E only distantly, contains 29 *Streptococcus* species (see Fig. S5C).

The phylum *Bacteroidetes* is represented by 72 species, which are found on three divergent clusters, K, N, and M, on the NanA tree (Fig. 5; also Fig. S6 in the supplemental material). Cluster K is a highly divergent group with long-branch lengths and is comprised of members of the phyla *Planctomycetes* and *Bacteroidetes*. Branching divergently from this group in cluster L are representatives of the phylum *Verrucomicrobia* (see Fig. S6A in the supple-

mental material). Cluster M consists of closely related NanA alleles from *Bacteroides*, *Prevotella*, and *Tannerella* species, many of which are human gut commensals, and all contain a NanT-type transporter (see Fig. S6B). *Tannerella forsythiae* and *Bacteroides fragilis* were shown to utilize sialic acid as a sole carbon source (63, 64).

The phylum *Actinobacteria* is represented by 83 species, the majority of which are present on the most divergent branches of the NanA tree, clusters Q, R, S, and T (Fig. 5; also Fig. S7 in the supplemental material). These clusters present mainly *Corynebacterium*, *Arthrobacter*, and *Bifidobacterium* species; nested within cluster S was a NanA from *Gardnerella vaginalis*, which suggests that it may have been acquired from a *Bifidobacterium* (see Fig. S7A in the supplemental material). All members of clusters Q, R, S, and T had an ABC type putative sialic acid transporter. Experimental evidence for sialic acid utilization was shown for *Gardnerella vaginalis*, *Corynebacterium diphtheriae*, *C. glutamicum*, and *Bifidobacterium breve* (65–68).

Conclusions. The adaptability to new nutritional sources is crucial for a species to switch between different environments. Adaptation to host environments requires the ability to switch from nutrient-poor to nutrient-rich and from free-living to host-associated conditions, including competing with the resident microbiota for both space and nutrients. Some of the major nutrient capabilities of *V. cholerae* are mucus associated, and pathogenic *V. cholerae* bacteria specifically scavenge and utilize sialic acids to allow niche expansion. Similarly, a recent study has proposed that the ability of *Salmonella enterica* and *Clostridium difficile* to colonize antibiotic-treated mice is due to the disruption of intestinal microbiota that allows these species to scavenge and utilize host sialic acids (69). It is of interest that, although the members of the TRAP family of transporters have a high affinity for sialic acid, their distribution is limited, being confined mainly to the genus *Vibrio* and the family *Pasteurellaceae*. Among other Gram-negative bacteria, NanT is the predominant sialic acid transporter, whereas the predominant types in Gram-positive bacteria are the SSS and ABC transporters. As the transporter distribution and phylogenetic tree suggest, sialic acid catabolism has evolved multiple times and is an ancestral trait in some species and newly acquired in others.

MATERIALS AND METHODS

Bacterial strains, plasmids, and growth conditions. All bacterial strains and plasmids used in this study are listed in Table S1 in the supplemental material. All *V. cholerae* strains were grown aerobically with aeration (225 rpm) at 37°C in Luria Bertani (LB) broth (Fisher Scientific, Waltham, MA) with a 1% NaCl concentration. For growth analysis on intestinal mucus, mucus sugars, and sialic acid derivatives as sole carbon sources, M9 minimal medium was supplemented with 0.02 mM MgSO₄ and 0.1 mM CaCl₂. M9 minimal medium supplemented with intestinal mucus (M9M) (30 µg/ml) or glucose (M9G) was used or *N*-acetylglucosamine, D-ribose, D-mannose, D-gluconate, or D-galactose (all 10 mM), or Neu5Ac (3 mM), 2-keto-3-deoxy-D-glycero-D-galactononic acid (KDN) (4 mM), or *N*-glycolylneuraminic acid (Neu5Gc) (3 mM) (Sigma-Aldrich) was added to M9 to serve as the carbon source. Antibiotics were used at the following concentrations: streptomycin (Sm) at 200 µg/ml, ampicillin (Amp) at 100 µg/ml, chloramphenicol (Cm) at 25 µg/ml, and kanamycin at 50 µg/ml.

Bacterial growth assays. Strains (diluted 1:100) were grown aerobically in LB broth at 37°C overnight and incubated for 3 h to reach mid-exponential-phase growth, and cells were pelleted by centrifugation, washed with M9 media, and resuspended in M9 media. The cultures were

then diluted 1:40 in M9 media supplemented with each carbon source and incubated at 37°C with intermittent shaking for 24 h. The optical density at 595 nm (OD_{595}) was taken hourly using a Tecan Sunrise microplate reader (Tecan Group, Ltd.) and Magellan plate reader software. High-throughput screens of carbon utilization abilities were performed using Biolog PM1 and PM2A plates as described above. Each experiment was performed with at least two biological replicates.

Complementation of the *E. coli* sialic acid transporter mutant. Genomic DNA from *V. cholerae* N16961 was used as a template for PCR amplification of *siaPQM* (VC1777-VC1779) transporter genes using primer pair VcsiaPQMF (TCTAGATGCTAAGCCATCAACATCTG) and VcsiaPQMR (GAGCTCTCAACTACTGTCCGTATCCG), which included XbaI and SacI restriction enzyme sites (underlined), respectively. The product was subcloned into pJET1.2 and transformed into *E. coli* DH5 α . Recombinant pJET1.2 plasmids were ligated into pBAD33 and were used to transform into *E. coli* JW3193, resulting in strain NDM3193C, which was used for the functional complementation assays described below. Strain NDM3193C was examined for growth in M9 media supplemented with arabinose (0.05% [wt/vol]) and with Neu5Ac, Neu5Gc, or KDN at 37°C. This construct was also used to complement NC1777 and demonstrate function complementation. Each strain and condition was tested using at least two biological replicates and in triplicate.

Mucus extraction. The intestinal mucus used in *in vitro* competition assays and growth analyses was isolated from mouse gastrointestinal tracts as previously described (70, 71). Briefly, mice gastrointestinal tracts were collected and the small intestine, cecum, and large intestine were first flushed with PBS followed by gentle scraping of the walls of the intestine to remove the mucus. The mucus was processed by homogenizing 200 mg of mucus in 5 ml of sterile PBS with streptomycin and centrifuged at 500 $\times$ g to pellet out tissue and fecal material. The protein content of the mucus was determined using a Bradford assay. Competition assays and growth analyses were performed in M9 media supplemented with mucus (final concentration of 30 μ g/ml of protein) (M9M).

***In vivo* and *in vitro* competition assays.** All experiments involving mice were approved by the University of Delaware Institutional Animal Care and Use Committee. Male C57BL/6 mice between 6 and 10 weeks of age were treated with streptomycin and inoculated as described previously (33, 70, 72). In short, the mice ($n = 10$) were fasted for 4 h before being orogastrically administered streptomycin and were then resupplied with food and water, supplemented with streptomycin (2.5 g/500 ml). At 20 h after antibiotic treatment, the mice were again fasted for 4 h and orogastrically inoculated with 100 μ l of bacterial suspension in PBS with a final concentration of 1×10^{10} CFU/ml. *Vibrio cholerae* strain SAM2338 (Δ lacZ) was used as a surrogate wild-type strain in competition experiments as it allows white/blue colony selection with the NC1777 strain (*AsiaM*) when grown on X-Gal LB plates.

For *in vitro* competition assays, an aliquot of the *in vivo* inoculum was used to inoculate M9G, M9M, or LB. Cultures were incubated at 37°C for 24 h, serially diluted and plated on LB X-Gal plates, and then incubated overnight at 37°C. CFU were counted as either blue (N16961 or NC1777) or white (SAM2338), and the ratio between the strains was determined. The competitive index (CI) was determined as follows: $CI = \text{ratio out}_{(NC1777 / SAM2338)} / \text{ratio in}_{(NC1777 / SAM2338)}$. A CI of <1 indicated that the wild-type strain outcompeted the mutant strain; a CI of >1 indicated the mutant strain outcompeted the wild-type strain. For the *in vivo* persistence competition assay, fecal pellets were collected (2 to 4 per animal) at daily time intervals from coinfected mice. The fecal pellets were weighed and then mechanically homogenized in 2 ml of sterile PBS, serially diluted and plated on LB Sm X-Gal agar, and incubated at 37°C overnight. The ratio of NC1777 colonies to SAM2338 colonies was termed the “ratio out” at each time point. After the last time point of the persistence assay, the mice were sacrificed and the entire gastrointestinal tract was harvested from each mouse and separated into samples of the small intestine, cecum, and large intestine. Each sample was homogenized in 4 ml of PBS,

serially diluted in PBS, plated on LB X-Gal plates, and incubated overnight at 37°C, and a blue/white colony screen was performed as described above.

Phylogenetic analysis. We queried using pBLAST with NanA of *V. cholerae* N16961 (VC1776) and *Staphylococcus aureus* N315 (SA0304) as seeds, and all completed and draft bacterial genomes (~7,400 genomes) were published on the National Center for Biotechnology Information (NCBI) website (73). In addition to a putative NanA, candidate species were determined by the presence of homologues to a sialic acid transporter, a putative *nanE* or *nanK* gene, or a gene encoding sialidase in proximity to the NanA-encoding gene on the genome. Sequences were aligned using ClustalW (see Table S2 in the supplemental material) (74). Aligned sequences were manually checked, and phylogenetic analysis was conducted with MEGA6 using the neighbor-joining (NJ) tree-building method and *p* distance and Poisson correction models for estimating evolutionary distances (75, 76).

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <http://mbio.asm.org/lookup/suppl/doi:10.1128/mBio.02237-15/-/DCSupplemental>.

Figure S1, TIF file, 1.9 MB.

Figure S2, TIF file, 1.9 MB.

Figure S3, TIF file, 0.2 MB.

Figure S4, TIF file, 0.1 MB.

Figure S5, TIF file, 0.2 MB.

Figure S6, TIF file, 0.1 MB.

Figure S7, TIF file, 0.2 MB.

Table S1, DOCX file, 0.03 MB.

ACKNOWLEDGMENTS

We gratefully acknowledge members of the Boyd Group for reviewing and giving constructive feedback on the manuscript.

This research was supported in part by a National Science Foundation CAREER award, DEB-0844409, to E.F.B. N.D.M. and J.-B.L. were supported in part by the Chemistry Biology Interface graduate program at the University of Delaware.

FUNDING INFORMATION

This work was funded by National Science Foundation (NSF) (DEB-0844409).

This research was supported in part by a National Science Foundation CAREER award DEB-0844409 to EFB

REFERENCES

- Ramamurthy T, Garg S, Sharma R, Bhattacharya SK, Balakrish Nair G, Shimada T, Takeda T, Karasawa T, Kurazano H, Pal A, Takeda Y. 1993. Emergence of novel strain of *Vibrio cholerae* with epidemic potential in southern and eastern India. *Lancet* 341:703–704. [http://dx.doi.org/10.1016/0140-6736\(93\)90480-5](http://dx.doi.org/10.1016/0140-6736(93)90480-5).
- Faruque SM, Sack DA, Sack RB, Colwell RR, Takeda Y, Nair GB. 2003. Emergence and evolution of *Vibrio cholerae* O139. *Proc Natl Acad Sci U S A* 100:1304–1309. <http://dx.doi.org/10.1073/pnas.0337468100>.
- Sack DA, Sack RB, Nair GB, Siddique AK. 2004. Cholera. *Lancet* 363: 223–233. [http://dx.doi.org/10.1016/S0140-6736\(03\)15328-7](http://dx.doi.org/10.1016/S0140-6736(03)15328-7).
- Waldor MK, Mekalanos JJ. 1994. Emergence of a new cholera pandemic: molecular analysis of virulence determinants in *Vibrio cholerae* O139 and development of a live vaccine prototype. *J Infect Dis* 170:278–283. <http://dx.doi.org/10.1093/infdis/170.2.278>.
- Jermyn WS, Boyd EF. 2002. Characterization of a novel *Vibrio* pathogenicity island (VPI-2) encoding neuraminidase (*nanH*) among toxigenic *Vibrio cholerae* isolates. *Microbiology* 148:3681–3693. <http://dx.doi.org/10.1099/002221287-148-11-3681>.
- Kovacikova G, Lin W, Skorupski K. 2005. Dual regulation of genes involved in acetoin biosynthesis and motility/biofilm formation by the virulence activator AphA and the acetate-responsive LysR-type regulator AlsR in *Vibrio cholerae*. *Mol Microbiol* 57:420–433. <http://dx.doi.org/10.1111/j.1365-2958.2005.04700.x>.
- Yoon SS, Mekalanos JJ. 2006. 2,3-Butanediol synthesis and the emer-

- gence of the *Vibrio cholerae* El Tor biotype. *Infect Immun* 74:6547–6556. <http://dx.doi.org/10.1128/IAI.00695-06>.
8. Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R, Gordon JI. 2007. The human microbiome project. *Nature* 449:804–810. <http://dx.doi.org/10.1038/nature06244>.
 9. Ley RE, Knight R, Gordon JI. 2007. The human microbiome: eliminating the biomedical/environmental dichotomy in microbial ecology. *Environ Microbiol* 9:3–4. http://dx.doi.org/10.1111/j.1462-2920.2006.01222_3.x.
 10. Gill SR, Pop M, Deboy RT, Eckburg PB, Turnbaugh PJ, Samuel BS, Gordon JI, Relman DA, Fraser-Liggett CM, Nelson KE. 2006. Metagenomic analysis of the human distal gut microbiome. *Science* 312:1355–1359. <http://dx.doi.org/10.1126/science.1124234>.
 11. Freter R. 1962. *In vivo* and *in vitro* antagonism of intestinal bacteria against *Shigella flexneri*. II. The inhibitory mechanism. *J Infect Dis* 110:38–46. <http://dx.doi.org/10.1093/infdis/110.1.38>.
 12. Stecher B, Hardt WD. 2011. Mechanisms controlling pathogen colonization of the gut. *Curr Opin Microbiol* 14:82–91. <http://dx.doi.org/10.1016/j.mib.2010.10.003>.
 13. Leatham MP, Banerjee S, Autieri SM, Mercado-Lubo R, Conway T, Cohen PS. 2009. Precolonized human commensal *Escherichia coli* strains serve as a barrier to *E. coli* O157:H7 growth in the streptomycin-treated mouse intestine. *Infect Immun* 77:2876–2886. <http://dx.doi.org/10.1128/IAI.00059-09>.
 14. Fabich AJ, Jones SA, Chowdhury FZ, Cernosek A, Anderson A, Smalley D, McHargue JW, Hightower GA, Smith JT, Autieri SM, Leatham MP, Lins JJ, Allen RL, Laux DC, Cohen PS, Conway T. 2008. Comparison of carbon nutrition for pathogenic and commensal *Escherichia coli* strains in the mouse intestine. *Infect Immun* 76:1143–1152. <http://dx.doi.org/10.1128/IAI.01386-07>.
 15. Bertin Y, Chaucheyras-Durand F, Robbe-Masselot C, Durand A, de la Foye A, Harel J, Cohen PS, Conway T, Forano E, Martin C. 2013. Carbohydrate utilization by enterohaemorrhagic *Escherichia coli* O157:H7 in bovine intestinal content. *Environ Microbiol* 15:610–622. <http://dx.doi.org/10.1111/1462-2920.12019>.
 16. Corfield AP, Carroll D, Myerscough N, Probert CS. 2001. Mucins in the gastrointestinal tract in health and disease. *Front Biosci* 6:D1321–D1357. <http://dx.doi.org/10.2741/Corfield>.
 17. Martens EC, Chiang HC, Gordon JI. 2008. Mucosal glycan foraging enhances fitness and transmission of a saccharolytic human gut bacterial symbiont. *Cell Host Microbe* 4:447–457. <http://dx.doi.org/10.1016/j.chom.2008.09.007>.
 18. Marcobal A, Southwick AM, Earle KA, Sonnenburg JL. 2013. A refined palate: bacterial consumption of host glycans in the gut. *Glycobiology* 23:1038–1046. <http://dx.doi.org/10.1093/glycob/cwt040>.
 19. Manzi AE, Norgard-Sumnicht K, Argade S, Marth JD, van Halbeek H, Varki A. 2000. Exploring the glycan repertoire of genetically modified mice by isolation and profiling of the major glycan classes and nano-NMR analysis of glycan mixtures. *Glycobiology* 10:669–689. <http://dx.doi.org/10.1093/glycob/10.7.669>.
 20. Bansil R, Turner BS. 2006. Mucin structure, aggregation, physiological functions and biomedical applications *Curr Opin Colloid Interface Sci* 11:164–170. <http://dx.doi.org/10.1016/j.cocis.2005.11.001>.
 21. Jermyn WS, Boyd EF. 2005. Molecular evolution of *Vibrio* pathogenicity island-2 (VPI-2): mosaic structure among *Vibrio cholerae* and *Vibrio mimicus* natural isolates. *Microbiology* 151:311–322. <http://dx.doi.org/10.1099/mic.0.27621-0>.
 22. Almagro-Moreno S, Boyd EF. 2009. Sialic acid catabolism confers a competitive advantage to pathogenic *Vibrio cholerae* in the mouse intestine. *Infect Immun* 77:3807–3816. <http://dx.doi.org/10.1128/IAI.00279-09>.
 23. Chowdhury N, Norris J, McAlister E, Lau SY, Thomas GH, Boyd EF. 2012. The VC1777-VC1779 proteins are members of a sialic acid-specific subfamily of TRAP transporters (SiaPQM) and constitute the sole route of sialic acid uptake in the human pathogen *Vibrio cholerae*. *Microbiology* 158:2158–2167. <http://dx.doi.org/10.1099/mic.0.059659-0>.
 24. Vimr E, Lichtensteiger C, Steenbergen S. 2000. Sialic acid metabolism's dual function in *Haemophilus influenzae*. *Mol Microbiol* 36:1113–1123. <http://dx.doi.org/10.1046/j.1365-2958.2000.01925.x>.
 25. Severi E, Hood DW, Thomas GH. 2007. Sialic acid utilization by bacterial pathogens. *Microbiology* 153:2817–2822. <http://dx.doi.org/10.1099/mic.0.2007/009480-0>.
 26. Almagro-Moreno S, Boyd EF. 2009. Insights into the evolution of sialic acid catabolism among bacteria. *BMC Evol Biol* 9:118. <http://dx.doi.org/10.1186/1471-2148-9-118>.
 27. Fischer M, Zhang QY, Hubbard RE, Thomas GH. 2010. Caught in a TRAP: substrate-binding proteins in secondary transport. *Trends Microbiol* 18:471–478. <http://dx.doi.org/10.1016/j.tim.2010.06.009>.
 28. Haines-Menges BL, Whitaker WB, Lubin JB, Boyd EF. 2015. Host sialic acids: a delicacy for the pathogen with discerning taste. *Microbiol Spectr* 3(4). <http://dx.doi.org/10.1128/microbiolspec.MBP-0005-2014>.
 29. Varki A. 2001. Loss of N-glycolylneuraminic acid in humans: mechanisms, consequences, and implications for hominid evolution. *Am J Phys Anthropol* 33(Suppl):54–69.
 30. Chou HH, Takematsu H, Diaz S, Iber J, Nickerson E, Wright KL, Muchmore EA, Nelson DL, Warren ST, Varki A. 1998. A mutation in human CMP-sialic acid hydroxylase occurred after the Homo-Pan divergence. *Proc Natl Acad Sci U S A* 95:11751–11756. <http://dx.doi.org/10.1073/pnas.95.20.11751>.
 31. Lubin JB, Kingston JJ, Chowdhury N, Boyd EF. 2012. Sialic acid catabolism and transport gene clusters are lineage specific in *Vibrio vulnificus*. *Appl Environ Microbiol* 78:3407–3415. <http://dx.doi.org/10.1128/AEM.07395-11>.
 32. Hopkins AP, Hawkhead JA, Thomas GH. 2013. Transport and catabolism of the sialic acids N-glycolylneuraminic acid and 3-keto-3-deoxy-D-glycero-D-galactononic acid by *Escherichia coli* K-12. *FEMS Microbiol Lett* 347:14–22. <http://dx.doi.org/10.1111/1574-6968.12213>.
 33. Nygren E, Li BL, Holmgren J, Attridge SR. 2009. Establishment of an adult mouse model for direct evaluation of the efficacy of vaccines against *Vibrio cholerae*. *Infect Immun* 77:3475–3484. <http://dx.doi.org/10.1128/IAI.01197-08>.
 34. Bohnhoff M, Miller CP. 1962. Enhanced susceptibility to salmonella infection in streptomycin-treated mice. *J Infect Dis* 111:117–127. <http://dx.doi.org/10.1093/infdis/111.2.117>.
 35. Miller CP, Bohnhoff M. 1963. Changes in the mouse's enteric microflora associated with enhanced susceptibility to salmonella infection following streptomycin treatment. *J Infect Dis* 113:59–66. <http://dx.doi.org/10.1093/infdis/113.1.59>.
 36. Miller CP, Bohnhoff M, Rifkind D. 1956–1957. The effect of an antibiotic on the susceptibility of the mouse's intestinal tract to salmonella infection. *Trans Am Clin Climatol Assoc* 68:51–55; discussion 55–58.
 37. Barthel M, Hapfelmeier S, Quintanilla-Martínez L, Kremer M, Rohde M, Högardt M, Pfeffer K, Rüssmann H, Hardt WD. 2003. Pretreatment of mice with streptomycin provides a *Salmonella enterica* serovar Typhimurium colitis model that allows analysis of both pathogen and host. *Infect Immun* 71:2839–2858. <http://dx.doi.org/10.1128/IAI.71.5.2839-2858.2003>.
 38. Myhal ML, Laux DC, Cohen PS. 1982. Relative colonizing abilities of human fecal and K 12 strains of *Escherichia coli* in the large intestines of streptomycin-treated mice. *Eur J Clin Microbiol* 1:186–192. <http://dx.doi.org/10.1007/BF02019621>.
 39. Wadolkowski EA, Laux DC, Cohen PS. 1988. Colonization of the streptomycin-treated mouse large intestine by a human fecal *Escherichia coli* strain: role of adhesion to mucosal receptors. *Infect Immun* 56:1036–1043.
 40. McCormick BA, Laux DC, Cohen PS. 1990. Neither motility nor chemotaxis plays a role in the ability of *Escherichia coli* F-18 to colonize the streptomycin-treated mouse large intestine. *Infect Immun* 58:2957–2961.
 41. Sweeney NJ, Klemm P, McCormick BA, Møller-Nielsen E, Utley M, Schembri MA, Laux DC, Cohen PS. 1996. The *Escherichia coli* K-12 gntP gene allows *E. coli* F-18 to occupy a distinct nutritional niche in the streptomycin-treated mouse large intestine. *Infect Immun* 64:3497–3503.
 42. Sweeney NJ, Laux DC, Cohen PS. 1996. *Escherichia coli* F-18 and *E. coli* K-12 eda mutants do not colonize the streptomycin-treated mouse large intestine. *Infect Immun* 64:3504–3511.
 43. Sekirov I, Tam NM, Jogova M, Robertson ML, Li Y, Lupp C, Finlay BB. 2008. Antibiotic-induced perturbations of the intestinal microbiota alter host susceptibility to enteric infection. *Infect Immun* 76:4726–4736. <http://dx.doi.org/10.1128/IAI.00319-08>.
 44. Jones SA, Chowdhury FZ, Fabich AJ, Anderson A, Schreiner DM, House AL, Autieri SM, Leatham MP, Lins JJ, Jørgensen M, Cohen PS, Conway T. 2007. Respiration of *Escherichia coli* in the mouse intestine. *Infect Immun* 75:4891–4899. <http://dx.doi.org/10.1128/IAI.00484-07>.
 45. Leatham-Jensen MP, Frimodt-Møller J, Adediran J, Mokszycki ME, Banner ME, Caughron JE, Krogfelt KA, Conway T, Cohen PS. 2012. The streptomycin-treated mouse intestine selects *Escherichia coli* envZ mis-sense mutants that interact with dense and diverse intestinal microbiota. *Infect Immun* 80:1716–1727. <http://dx.doi.org/10.1128/IAI.06193-11>.

46. Stern AM, Hay AJ, Liu Z, Desland FA, Zhang J, Zhong Z, Zhu J. 2012. The NorR regulon is critical for *Vibrio cholerae* resistance to nitric oxide and sustained colonization of the intestines. *mBio* 3(2):e00013-12. <http://dx.doi.org/10.1128/mBio.00013-12>.
47. Maltby R, Leatham-Jensen MP, Gibson T, Cohen PS, Conway T. 2013. Nutritional basis for colonization resistance by human commensal *Escherichia coli* strains HS and Nissle 1917 against *E. coli* O157:H7 in the mouse intestine. *PLoS One* 8:e53957. <http://dx.doi.org/10.1371/journal.pone.0053957>.
48. Spees AM, Wangdi T, Lopez CA, Kingsbury DD, Xavier MN, Winter SE, Tsolis RM, Bäumler AJ. 2013. Streptomycin-induced inflammation enhances *Escherichia coli* gut colonization through nitrate respiration. *mBio* 4(4):e00430-13. <http://dx.doi.org/10.1128/mBio.00430-13>.
49. Meador JP, Caldwell ME, Cohen PS, Conway T. 2014. *Escherichia coli* pathotypes occupy distinct niches in the mouse intestine. *Infect Immun* 82:1931–1938. <http://dx.doi.org/10.1128/IAI.01435-13>.
50. Schinner SA, Mokszycki ME, Adediran J, Leatham-Jensen M, Conway T, Cohen PS. 2015. *Escherichia coli* EDL933 Requires gluconeogenic nutrients to successfully colonize the intestines of streptomycin-treated mice pre-colonized with *E. coli* Nissle 1917. *Infect Immun* 83:1983–1991. <http://dx.doi.org/10.1128/IAI.02943-14>.
51. Gonin S, Arnoux P, Pierru B, Lavergne J, Alonso B, Sabaty M, Pignol D. 2007. Crystal structures of an extracytoplasmic solute receptor from a TRAP transporter in its open and closed forms reveal a helix-swapped dimer requiring a cation for alpha-keto acid binding. *BMC Struct Biol* 7:11. <http://dx.doi.org/10.1186/1472-6807-7-11>.
52. Mulligan C, Leech AP, Kelly DJ, Thomas GH. 2012. The membrane proteins SiaQ and SiaM form an essential stoichiometric complex in the sialic acid tripartite ATP-independent periplasmic (TRAP) transporter SiaPQM (VC1777-1779) from *Vibrio cholerae*. *J Biol Chem* 287:3598–3608. <http://dx.doi.org/10.1074/jbc.M111.281030>.
53. Post DM, Mungur R, Gibson BW, Munson RS, Jr. 2005. Identification of a novel sialic acid transporter in *Haemophilus ducreyi*. *Infect Immun* 73:6727–6735. <http://dx.doi.org/10.1128/IAI.73.10.6727-6735.2005>.
54. Steenbergen SM, Lichtensteiger CA, Caughlan R, Garfinkle J, Fuller TE, Vimr ER. 2005. Sialic acid metabolism and systemic Pasteurellosis. *Infect Immun* 73:1284–1294. <http://dx.doi.org/10.1128/IAI.73.3.1284-1294.2005>.
55. Allen S, Zaleski A, Johnston JW, Gibson BW, Apicella MA. 2005. Novel sialic acid transporter of *Haemophilus influenzae*. *Infect Immun* 73:5291–5300. <http://dx.doi.org/10.1128/IAI.73.9.5291-5300.2005>.
56. Severi E, Randle G, Kivlin P, Whitfield K, Young R, Moxon R, Kelly D, Hood D, Thomas GH. 2005. Sialic acid transport in *Haemophilus influenzae* is essential for lipopolysaccharide sialylation and serum resistance and is dependent on a novel tripartite ATP-independent periplasmic transporter. *Mol Microbiol* 58:1173–1185. <http://dx.doi.org/10.1111/j.1365-2958.2005.04901.x>.
57. Johnston JW, Zaleski A, Allen S, Mootz JM, Armbruster D, Gibson BW, Apicella MA, Munson RS, Jr. 2007. Regulation of sialic acid transport and catabolism in *Haemophilus influenzae*. *Mol Microbiol* 66:26–39. <http://dx.doi.org/10.1111/j.1365-2958.2007.05890.x>.
58. Vimr ER, Troy FA. 1985. Identification of an inducible catabolic system for sialic acids (*nan*) in *Escherichia coli*. *J Bacteriol* 164:845–853.
59. Severi E, Hosie AH, Hawkhead JA, Thomas GH. 2010. Characterization of a novel sialic acid transporter of the sodium solute symporter (SSS) family and in vivo comparison with known bacterial sialic acid transporters. *FEMS Microbiol Lett* 304:47–54. <http://dx.doi.org/10.1111/j.1574-6968.2009.01881.x>.
60. Joseph S, Hariri S, Masood N, Forsythe S. 2013. Sialic acid utilization by *Cronobacter sakazakii*. *Microb Inform Exp* 3:3. <http://dx.doi.org/10.1186/2042-5783-3-3>.
61. Molenaar D, Bringel F, Schuren FH, de Vos WM, Siezen RJ, Kleerebezem M. 2005. Exploring *Lactobacillus plantarum* genome diversity by using microarrays. *J Bacteriol* 187:6119–6127. <http://dx.doi.org/10.1128/JB.187.17.6119-6127.2005>.
62. Marco ML, Peters TH, Bongers RS, Molenaar D, van Hemert S, Sonnenburg JL, Gordon JI, Kleerebezem M. 2009. Lifestyle of *Lactobacillus plantarum* in the mouse caecum. *Environ Microbiol* 11:2747–2757. <http://dx.doi.org/10.1111/j.1462-2920.2009.02001.x>.
63. Roy S, Douglas CW, Stafford GP. 2010. A novel sialic acid utilization and uptake system in the periodontal pathogen *Tannerella forsythiae*. *J Bacteriol* 192:2285–2293. <http://dx.doi.org/10.1128/JB.00079-10>.
64. Brigham C, Caughlan R, Gallegos R, Dallas MB, Godoy VG, Malamy MH. 2009. Sialic acid (N-acetyl neuraminic acid or NANA) utilization by *Bacteroides fragilis* requires a novel N-acetyl mannosamine (manNAc) epimerase. *J Bacteriol* 191:3629–3638. <http://dx.doi.org/10.1128/JB.00811-08>.
65. Egan M, O'Connell Motherway M, Ventura M, van Sinderen D. 2014. Metabolism of sialic acid by *Bifidobacterium breve* UCC2003. *Appl Environ Microbiol* 80:4414–4426. <http://dx.doi.org/10.1128/AEM.01114-14>.
66. Gilbert NM, Lewis WG, Lewis AL. 2013. Clinical features of bacterial vaginosis in a murine model of vaginal infection with *Gardnerella vaginalis*. *PLoS One* 8:e59539. <http://dx.doi.org/10.1371/journal.pone.0059539>.
67. Lewis WG, Robinson LS, Gilbert NM, Perry JC, Lewis AL. 2013. Degradation, foraging, and depletion of mucus sialoglycans by the vagina-adapted *Actinobacterium Gardnerella vaginalis*. *J Biol Chem* 288:12067–12079. <http://dx.doi.org/10.1074/jbc.M113.453654>.
68. Arden SB, Chang WH, Barksdale L. 1972. Distribution of neuraminidase and n-acetylneuraminase activities among corynebacteria, mycobacteria, and nocardias. *J Bacteriol* 112:1206–1212.
69. Ng KM, Ferreyra JA, Higginbottom SK, Lynch JB, Kashyap PC, Gopinath S, Naidu N, Choudhury B, Weimer BC, Monack DM, Sonnenburg JL. 2013. Microbiota-liberated host sugars facilitate post-antibiotic expansion of enteric pathogens. *Nature* 502:96–99. <http://dx.doi.org/10.1038/nature12503>.
70. Whitaker WB, Richards GP, Boyd EF. 2014. Loss of sigma factor RpoN increases intestinal colonization of *Vibrio parahaemolyticus* in an adult mouse model. *Infect Immun* 82:544–556. <http://dx.doi.org/10.1128/IAI.01210-13>.
71. Haines-Menges B, Whitaker WB, Boyd EF. 2014. Alternative sigma factor RpoE is important for *Vibrio parahaemolyticus* cell envelope stress response and intestinal colonization. *Infect Immun* 82:3667–3677. <http://dx.doi.org/10.1128/IAI.01854-14>.
72. Whitaker WB, Parent MA, Boyd A, Richards GP, Boyd EF. 2012. The *Vibrio parahaemolyticus* ToxRS regulator is required for stress tolerance and colonization in a novel orogastric streptomycin-induced adult murine model. *Infect Immun* 80:1834–1845. <http://dx.doi.org/10.1128/IAI.06284-11>.
73. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and psi-blast: a new generation of protein database search programs. *Nucleic Acids Res* 25:3389–3402. <http://dx.doi.org/10.1093/nar/25.17.3389>.
74. Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22:4673–4680. <http://dx.doi.org/10.1093/nar/22.22.4673>.
75. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30:2725–2729. <http://dx.doi.org/10.1093/molbev/mst197>.
76. Saitou N, Nei M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425.
77. Zuckerkandl E, Pauling L. 1965. Evolutionary divergence and convergence in proteins, p 97–166. In Bryson V, Vogel HJ (ed), *Evolving Genes and Proteins*. Academic Press, New York, NY.